

in cancer patients who have been treated with exogenous interferon gamma or interleukin 2.⁴ The latter is a cytokine which produces significant eosinophilia.⁵ Thus, in all likelihood the abnormal tryptophan metabolism seen in EMS patients is a consequence of the immune activation in this disease rather than the cause of it. Indeed, one might reason that an important factor in the pathogenesis of EMS may be an excess, rather than a deficit of interferon gamma. The observed active catabolism of tryptophan by the IDO pathway may well lead to a deficiency of tryptophan and serotonin. If serotonin does play a role in down-regulating interferon production as suggested,⁶ this deficit of serotonin may lead to an unchecked increase in interferon gamma and result in excessive activation of macrophages and eosinophils as seen in EMS.

In addition, although tryptophan metabolism may play a role in the pathogenesis of EMS, epidemiologic studies point increasingly to a contaminant as the principal causal factor.⁷⁻⁹ The recent identification of the chemical structure of one such contaminant¹⁰ may soon lead to animal and in vitro models which will permit dissection of this most interesting syndrome.

PHILLIP A. HERTZMAN, MD
Los Alamos Medical Center
Los Alamos, NM 87544
RAYMOND R. BROWN, PhD
University of Wisconsin
Clinical Cancer Center
Madison, WI 53706

REFERENCES

1. Criswell LA, Sack KE: Tryptophan-induced eosinophilia-myalgia syndrome. *West J Med* 1990 Sep; 153:269-274
2. Silver RM, Heyes MP, Maize JC, Quearry B, Vionnet-Fuasset M, Sternberg EM: Scleroderma, fasciitis, and eosinophilia associated with the ingestion of tryptophan. *N Engl J Med* 1990; 322:874-881
3. Byrne GI, Lehmann LK, Kirschbaum JG, Borden EC, Lee CM, Brown RR: Induction of tryptophan degradation in vitro and in vivo: A gamma-interferon-stimulated activity. *J Interferon Res* 1986; 6:389-396
4. Brown RR, Lee CM, Kohler PC, Hank JA, Storer BE, Sondel PM: Altered tryptophan and neopterin metabolism in cancer patients treated with recombinant interleukin-2. *Cancer Res* 1989; 49:4941-4944
5. Sedgwick JB, Frick WE, Sondel PM, Hank JA, Borden E, Busse WW: The appearance of hypodense eosinophils during interleukin-2 treatment. *J Allergy Clin Immunol* 1990; 85:557-566
6. Artz ES, Fernandez-Castelo S, Finocchiaro L, et al: Immunomodulation by indoleamines: Serotonin and melatonin action on DNA and interferon-gamma synthesis by human peripheral blood mononuclear cells. *J Clin Immunol* 1988; 8:513-520
7. Belongia EA, Hedberg CW, Gleich GJ, et al: An investigation of the cause of the eosinophilia-myalgia syndrome associated with tryptophan use. *N Engl J Med* 1990; 323:357-365
8. Slutsker L, Hoesly FC, Miller L, Williams P, Watson JC, Fleming DW: Eosinophilia-myalgia syndrome associated with exposure to tryptophan from a single manufacturer. *JAMA* 1990; 264:213-217

9. Centers for Disease Control: Update: Analysis of L-tryptophan for the etiology of eosinophilia-myalgia syndrome. *MMWR* 1990; 39:789-790

10. Sakimoto K: The cause of the eosinophilia-myalgia syndrome associated with tryptophan use. *N Engl J Med* 1990; 323:992-993

* * *

Drs Criswell and Sack Respond

TO THE EDITOR: We are pleased to respond to the thoughtful comments of Drs Hertzman and Brown. No one has measured interferon-gamma (IFN- γ) levels in patients with the eosinophilia-myalgia syndrome (EMS); thus, the finding by Silver and co-workers of elevated serum levels of L-kynurenine and quinolinic acid in such persons is unexplained.¹

Interferon gamma can indeed stimulate indoleamine-2,3-dioxygenase (IDO), the rate limiting enzyme of the kynurenine pathway of L-tryptophan metabolism, but other substances such as endotoxin can also induce this enzyme. Furthermore, L-tryptophan influences the ability of IFN- γ to induce IDO.

Products of the indoleamine pathway of L-tryptophan metabolism, serotonin and melatonin, inhibit the production of IFN- γ . This may be important in the pathogenesis of EMS because IFN- γ is known to inhibit synthesis of collagen, the production of interleukins 4 and 5, and the production of IgE by B lymphocytes.²⁻⁵

It is likely that most, if not all, of the current cases of EMS are associated with a contaminant generated during the manufacturing of L-tryptophan. The manner in which L-tryptophan or a contaminant induces EMS is still open to speculation, however.

LINDSEY A. CRISWELL, MD
KENNETH E. SACK, MD
Division of Rheumatology
University of California, San Francisco,
School of Medicine
400 Parnassus Ave, A-540
San Francisco, CA 94143-0362

REFERENCES

1. Silver RM, Heyes MP, Maize JC, Quearry B, Vionnet-Fuasset M, Sternberg EM: Scleroderma, fasciitis, and eosinophilia associated with the ingestion of tryptophan. *N Engl J Med* 1990; 322:874-881
2. Heckmann M, Aumailley M, Hatamochi A, Chu ML, Timpl R, Krieg T: Down-regulation of alpha-3 (VI) chain expression by γ interferon decreases synthesis and deposition of collagen type VI. *Eur J Biochem* 1989; 182:719-726
3. Coffman RL, Seymour BWP, Lebman DA, et al: The role of helper T cell products in mouse B cell differentiation and isotype regulation. *Immunol Rev* 1988; 102:5-28
4. Mossmann TR, Coffman RL: TH1 and TH2 cells: Different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 1989; 7:145-173
5. Pène J, Rousset F, Brière F, et al: IgE production by normal human lymphocytes is induced by interleukin 4 and suppressed by interferons gamma and alpha and prostaglandin E₂. *Proc Natl Acad Sci USA* 1988; 85:6880-6884